

Phone 201.854.4645

Fax 201.854.3203

235 60th Street

West New York, NJ 07093

INSTRUCTIONS FOR COLONOSCOPY

THE DAY BEFORE THE PROCEDURE YOU ARE TO BE ON LIQUID DIET ALL DAY WHICH CONSIST OF THE FOLLOWING:

- CHICKEN BROTH
- LIPTON TEA
- GATORADE (EXCEPT THE RED OR BLUE)
- JELLO (ONLY THE YELLOW COLOR)
- ONE CUP OF COFFEE
- LOTS OF WATER

****YOU SHOULD NOT DRINK ANYTHING THAT COTAINS RED COLORING, COCA COLA OR MILK (NOTHING SOLID) ****

Your procedure will be performed by Doctor: _____

Address: _____

Date of procedure: _____ Arrive at: _____ AM/PM

Comments: _____

SUPREP[®] BOWEL PREP KIT

(sodium sulfate, potassium sulfate and magnesium sulfate)
Oral Solution

(17.5g/3.13g/1.6g) per 6 ounces



On the day before your procedure...

What You CAN do:

- Do not eat or have clear liquids ONLY; please have nothing for dinner

What You CANNOT do:

- **DO NOT** drink milk
- **DO NOT** eat or drink anything colored red or purple
- **DO NOT** drink alcoholic beverages

Any of the following liquids are OK to drink:

- Water
- Strained fruit juices (without pulp) including apple, orange, white grape, or white cranberry
- Limeade or lemonade
- Coffee or tea (**DO NOT** use any dairy or non-dairy creamer)
- Chicken broth
- Gelatin desserts without added fruit or topping (**NO RED OR PURPLE**)

Note:

- SUPREP Bowel Prep Kit is indicated for cleansing of the colon as a preparation for colonoscopy in adults
- Be sure to tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. SUPREP Bowel Prep Kit may affect how other medicines work
- Medication taken by mouth may not be absorbed properly when taken within 1 hour before the start of each dose of SUPREP Bowel Prep Kit
- The most common adverse reactions after administration of SUPREP Bowel Prep Kit were overall discomfort, abdominal distention, abdominal pain, nausea, vomiting, and headache
- If you have any questions about taking SUPREP Bowel Prep Kit, call your doctor

To learn more about this product, please call **1-800-874-6756** or visit our Web site at **www.suprepkit.com**

Split-Dose (2-Day) Regimen

Begin Step 1 at _____ PM the evening before your procedure and proceed as shown below:


- On the evening before the procedure (or when your doctor tells you to begin) complete steps 1 through 4 using one (1) 6-ounce bottle before going to bed
- On the morning of your procedure, repeat steps 1 through 4 using the other 6-ounce bottle

STEP 1



Pour **ONE (1)** 6-ounce bottle of SUPREP liquid into the mixing container.

STEP 2




Add cool drinking water to the 16-ounce line on the container and mix.

STEP 3



Drink **ALL** the liquid in the container.

STEP 4



IMPORTANT
You **must** drink two (2) more 16-ounce containers of water over the next 1 hour.

Begin Step 1 at _____ AM the morning of your procedure and proceed as shown above:

NOTE: Dilute the solution concentrate as directed prior to use.

NOTE: You **must** finish drinking the final glass of water at least **6** hours, or as directed, before your colonoscopy.

Please read the Full Prescribing Information and Medication Guide in the kit.

Provided as an educational service from  **Braintree**
A PART OF SEBELA PHARMACEUTICALS[®]

©2019 Braintree. A part of Sebelo Pharmaceuticals

SU-12253A

April, 2019